

4. Approximation de fonctions

Exercice 4.1

a. $u_n = \frac{4}{n!}$ $A = \lim_{n \rightarrow \infty} \frac{\frac{4}{(n+1)!}}{\frac{4}{n!}} = \lim_{n \rightarrow \infty} \frac{4}{(n+1)!} \cdot \frac{n!}{4} = \lim_{n \rightarrow \infty} \frac{1}{n+1} = 0$

$\Rightarrow r = \frac{1}{0} = \infty \Rightarrow$ converge $\forall x$

b. $u_n = 2^n + 1$ $A = \lim_{n \rightarrow \infty} \frac{2^{n+1} + 1}{2^n + 1} \stackrel{\text{L'Hôpital}}{=} \lim_{n \rightarrow \infty} \frac{2^{n+1} \cdot \ln(2)}{2^n \cdot \ln(2)} = 2 \Rightarrow r = \frac{1}{2}$

Quand $x = \frac{1}{2}$: $\sum (2^n + 1) \cdot \left(\frac{1}{2}\right)^n = \sum \left(1 + \frac{1}{2^n}\right)$ diverge (critère de convergence)

Quand $x = -\frac{1}{2}$: $\sum (2^n + 1) \left(-\frac{1}{2}\right)^n = \sum \left(\frac{2^n}{(-2)^n} - \frac{1}{2^n}\right) = \sum \left((-1)^n - \frac{1}{2^n}\right)$ diverge

\Rightarrow converge si $-\frac{1}{2} < x < \frac{1}{2}$

c. $u_n = \frac{1}{n^n}$ $C = \lim_{n \rightarrow \infty} \frac{\sqrt[n]{\frac{1}{n^n}}}{\sqrt[n]{\frac{1}{n^n}}} = \lim_{n \rightarrow \infty} \frac{1}{n} = 0 \Rightarrow r = \infty$

\Rightarrow converge $\forall x$

d. $u_n = \frac{3}{(n+1)^n}$ $C = \lim_{n \rightarrow \infty} \frac{\sqrt[n]{\frac{3}{(n+1)^n}}}{\sqrt[n]{\frac{3}{(n+1)^n}}} = \lim_{n \rightarrow \infty} \frac{\sqrt[n]{3}}{n+1} = \frac{1}{\infty} = 0 \Rightarrow r = \infty$

\Rightarrow converge $\forall x$

e. $u_n = \frac{(-1)^{n+1}}{n}$ $C = \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+2}}{n+1}}{\frac{(-1)^{n+1}}{n}} \right| = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+2}}{(-1)^{n+1}} \cdot \frac{n}{n+1} \right| = 1 \Rightarrow r = 1$

Quand $x = -1$: $\sum \frac{(-1)^{n+1}}{n} (-1)^n = \sum \frac{(-1)^{2n+1}}{n} = -\sum \frac{1}{n}$ diverge

Quand $x = 1$: $\sum \frac{(-1)^{n+1}}{n} (1)^n = \sum \frac{(-1)^{n+1}}{n}$ converge (critère des séries alternées)

\Rightarrow converge si $-1 < x \leq 1$

Exercice 4.2

a. $\sum_{n \geq 1} n x^n \Rightarrow u_n = n \quad A = \lim_{n \rightarrow \infty} \left| \frac{n+1}{n} \right| = 1 \Rightarrow r = 1$

Quand $x = -1$: $\sum n(-1)^n$ diverge
 Quand $x = 1$: $\sum n \cdot 1^n = \sum n$ diverge } \Rightarrow converge si $-1 < x < 1$

b. $\sum_{n \geq 1} \frac{(-1)^{n+1}}{n} x^n \Rightarrow u_n = \frac{(-1)^{n+1}}{n}$ Voir ex 4.1 e...

c. $\sum_{n \geq 1} \frac{x^n}{n \cdot (n+1)} \Rightarrow u_n = \frac{1}{n(n+1)} \quad A = \lim_{n \rightarrow \infty} \left| \frac{\frac{1}{(n+2)(n+1)}}{\frac{1}{(n+1)n}} \right| = \lim_{n \rightarrow \infty} \frac{(n+1) \cdot n}{(n+2)(n+1)} = 1$
 $\Rightarrow r = 1$

Quand $x = -1$: $\sum \frac{(-1)^n}{n(n+1)}$ converge (th 3.2)
 Quand $x = 1$: $\sum \frac{1^n}{n(n+1)} = 1 \cdot (v.p. 24)$ } \Rightarrow converge si $-1 \leq x \leq 1$

d. $\sum_{n \geq 1} \frac{(-1)^{n+1} \cdot x^n}{n \cdot 5^n} \Rightarrow u_n = \frac{(-1)^{n+1}}{n \cdot 5^n}$

$A = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+2}}{(n+1)5^{n+1}} \cdot \frac{n \cdot 5^n}{(-1)^{n+1}} \right| = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+2}}{(-1)^{n+1}} \cdot \frac{5^n}{5^{n+1}} \cdot \frac{n}{n+1} \right| = \frac{1}{5} \Rightarrow r = 5$

Quand $x = -5$: $\sum \frac{(-1)^{n+1} \cdot (-5)^n}{n \cdot 5^n} = \sum \frac{(-1)^{n+1} \cdot (-1)^n \cdot 5^n}{n \cdot 5^n} = \sum \frac{-1}{n}$ diverge

Quand $x = 5$: $\sum \frac{(-1)^{n+1} \cdot 5^n}{n \cdot 5^n} = \sum \frac{(-1)^{n+1}}{n}$ converge (th 3.2)

\Rightarrow converge si $-5 < x \leq 5$

e. $\sum_{n \geq 1} \frac{x^{2n-2}}{n(n+1)(n+2)} \quad A = \lim_{n \rightarrow \infty} \left| \frac{x^{2(n+1)-2}}{(n+1)(n+2)(n+3)} \cdot \frac{n(n+1)(n+2)}{x^{2n-2}} \right| =$

méthode un peu différente mais le principe est le même

$= \lim_{n \rightarrow \infty} \left(\frac{n}{n+3} \right) \cdot \frac{x^{2n}}{x^{2n-2}} = x^2 \Rightarrow r = \frac{1}{x^2}$
 $\rightarrow 1 \cdot \frac{1}{x^2} = x^2$

\Rightarrow converge si $-1 < x < 1$ suite \rightarrow

Quand $x = -1$: $\sum_{n \geq 1} \frac{(-1)^{2n-2}}{n(n+1)(n+2)} = \sum_{n \geq 1} \frac{1}{n(n+1)(n+2)}$ }
 Quand $x = 1$: $\sum_{n \geq 1} \frac{1^{2n-2}}{n(n+1)(n+2)} = \sum_{n \geq 1} \frac{1}{n(n+1)(n+2)}$ }
 } $\left. \begin{array}{l} \text{converge} \\ \text{car } \sum \frac{1}{n(n+1)} \\ \text{converge (p.24)} \end{array} \right\}$

\Rightarrow converge si $-1 \leq x \leq 1$

f. $\sum_{n \geq 2} \left(\frac{x}{\ln(n)}\right)^n \Rightarrow u_n = \frac{1}{(\ln(n))^n}$ $\lim_{n \rightarrow \infty} \sqrt[n]{\frac{1}{(\ln(n))^n}} = \lim_{n \rightarrow \infty} \frac{1}{\ln(n)} = 0$
 $\Rightarrow r = \infty \Rightarrow$ converge $\forall x$

Exercice 4.3

a. $f(x) = \sin(x)$ $f'(x) = \cos(x)$ $f''(x) = -\sin(x)$ $f^{(3)}(x) = -\cos(x)$

$f(x) = \sin(0) + \frac{\cos(0)}{1!}x - \frac{\sin(0)}{2!}x^2 - \frac{\cos(0)}{3!}x^3 + \dots$
 $= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$

rayon de convergence : $A = \lim_{n \rightarrow +\infty} \left| \frac{(-1)^{n+1} x^{2n+3}}{(2n+3)!} \cdot \frac{(2n+1)!}{(-1)^n x^{2n+1}} \right|$
 $= \lim_{n \rightarrow +\infty} \left| \frac{-x^2}{(2n+3)(2n+2)} \right| \rightarrow 0 \cdot x^2$

converge si $0 \cdot x^2 < 1 \Rightarrow |x| < +\infty$

b. $f(x) = f'(x) = f''(x) = f^{(3)}(x) = \dots = e^x$

$f(x) = e^0 + e^0 x + \frac{e^0}{2!}x^2 + \frac{e^0}{3!}x^3 + \dots = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$

$A = \lim_{n \rightarrow +\infty} \left| \frac{x^{n+1}}{(n+1)!} \cdot \frac{n!}{x^n} \right| = \lim_{n \rightarrow +\infty} \left| \frac{x}{n+1} \right| \rightarrow 0 \cdot x$

converge si $0 \cdot x < 1 \Rightarrow |x| < +\infty$

$$c. f(x) = \ln(1+x) \quad f'(x) = \frac{1}{1+x} \quad f''(x) = \frac{-1}{(1+x)^2} \quad f^{(3)}(x) = \frac{2}{(1+x)^3}$$

$$f^{(4)}(x) = \frac{-6}{(1+x)^4} = \frac{-3!}{(1+x)^4}$$

(4)

$$f(x) = \ln(1) + \frac{1}{1+0}x - \frac{1}{2!} \frac{1}{(1+0)^2}x^2 + \frac{1}{3!} \frac{2}{(1+0)^3}x^3 - \frac{1}{4!} \frac{3!}{(1+0)^4}x^4$$

$$= 0 + x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

$$A = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+2} x^{n+1}}{n+1} \cdot \frac{n}{(-1)^{n+1} x^n} \right| = \lim_{n \rightarrow \infty} \left| -x \frac{n}{n+1} \right| = x$$

$\underbrace{\frac{n}{n+1}}_{\rightarrow 1}$

Converge si: $|x| < 1$

Exercice 4.4 voir corrigé

Exercice 4.5

$$a. f(x) = \ln(2x+5) \quad f'(x) = \frac{2}{2x+5} \quad f''(x) = \frac{-4}{(2x+5)^2}$$


$$f^{(3)}(x) = \frac{16}{(2x+5)^3} \quad f^{(4)}(x) = \frac{-96}{(2x+5)^4} \quad f^{(5)}(x) = \frac{768}{(2x+5)^5}$$

$$f(x) = \ln(5) + \frac{2}{5}x - \frac{2}{25}x^2 + \frac{8}{375}x^3 - \frac{4}{625}x^4 + \frac{32}{15 \cdot 625}x^5 + R_5(x)$$

$$f^{(6)}(x) = \frac{-7680}{(2x+5)^6}$$

$$|R_5(x)| \leq \frac{|x|^6}{6!} 7680 = \frac{32}{3} x^6$$


↑
 Π: plus grande valeur possible de $|f^{(6)}(x)|$
 dans l'intervalle $[-2; 2]$
 Ici, c'est quand $x = -2$



$b. f(x) = \frac{1}{x^2} + 4 \quad f'(x) = -\frac{2}{x^3} \quad f''(x) = \frac{6}{x^4} \quad f^{(3)}(x) = \frac{-24}{x^5} \quad f^{(4)}(x) = \frac{120}{x^6}$
 $f(-2) = \frac{17}{4} \quad f'(-2) = \frac{1}{4} \quad f''(-2) = \frac{3}{8} \quad f^{(3)}(-2) = \frac{3}{4} \quad f^{(4)}(-2) = \frac{15}{8}$

$f(x) = \frac{17}{4} + \frac{1}{4}(x+2) + \frac{3}{16}(x+2)^2 + \frac{1}{8}(x+2)^3 + \frac{5}{64}(x+2)^4 + R_4(x)$

$f^{(5)}(x) = \frac{-720}{x^7} \Rightarrow R_4(x) \leq \frac{|x+2|^5}{5!} \cdot 720 = 6|x+2|^5$
 $|f^{(5)}(-1)|$



Exercice 4.6

$f(x) = \ln(x) \quad f'(x) = \frac{1}{x} \quad f''(x) = -\frac{1}{x^2} \quad f^{(3)}(x) = \frac{2}{x^3} \quad f^{(4)}(x) = \frac{-6}{x^4} \quad f^{(5)}(x) = \frac{24}{x^5}$
 $f(x) = x-1 - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{3} - \frac{(x-1)^4}{4} + \frac{(x-1)^5}{5}$ (degré 5)



Exercise 4.7

$$a. f(x) = \frac{1}{4x} \quad f'(x) = \frac{-1}{4x^2} \quad f''(x) = \frac{1}{2x^3} \quad f^{(3)}(x) = -\frac{3}{2x^4} \quad f^{(4)}(x) = \frac{6}{x^5}$$

$$f(5) = \frac{1}{20} \quad f'(5) = \frac{-1}{100} \quad f''(5) = \frac{1}{250} \quad f^{(3)}(5) = -\frac{3}{1250}$$

$$f(x) = \frac{1}{20} - \frac{1}{100}(x-5) + \frac{1}{500}(x-5)^2 - \frac{1}{2500}(x-5)^3 + \dots$$

$$= \frac{1}{20} \sum_{k=0}^{\infty} (-1)^k \frac{(x-5)^k}{5^k}$$

$$b. f(x) = \cos(2x) \quad f'(x) = -2\sin(2x) \quad f''(x) = -4\cos(2x) \quad f^{(3)}(x) = 8\sin(2x)$$

$$f\left(\frac{\pi}{4}\right) = 0 \quad f'\left(\frac{\pi}{4}\right) = -2 \quad f''\left(\frac{\pi}{4}\right) = 0 \quad f^{(3)}\left(\frac{\pi}{4}\right) = 8$$

$$f^{(4)}(x) = 16\cos(2x) \quad f^{(5)}(x) = -32\sin(2x)$$

$$f^{(4)}\left(\frac{\pi}{4}\right) = 0 \quad f^{(5)}\left(\frac{\pi}{4}\right) = -32$$

$$f(x) = -2\left(x - \frac{\pi}{4}\right) + \frac{8}{3!}\left(x - \frac{\pi}{4}\right)^3 - \frac{32}{5!}\left(x - \frac{\pi}{4}\right)^5 + \dots$$

$$= \sum_{k=1}^{\infty} (-1)^k \frac{2^{2k-1}}{(2k-1)!} \left(x - \frac{\pi}{4}\right)^{2k-1}$$
